


What is claimed is


1. The fungicidal composition, which comprises:
 - (i) an agrochemical selected from the group consisting of methyl (2E)-3-methoxy-2-[2'-[[[3''-(1'''-fluoro-2'''-phenyl-1'''-ethenyloxy)phenyl] methylimino]oxy]methylphenyl]propenoate of formula (I); *N*-methyl (2E)-2-methoxyimino-2-[2'-[[[3''-(1'''-fluoro-2'''-phenyl-1'''-ethenyloxy)phenyl] methylimino]oxy]methylphenylacetamide of formula (II) and a mixture thereof, and
 - (ii) an adjuvant for enhancing the fungicidal efficacy, selected from the group consisting of: a nonionic surfactant selected from the group consisting of polyoxyethylene alkyl ether, polyoxyethylene alkyl ester, ethoxylated caster oil, polyoxyethylene-polyoxypropylene block copolymer and a mixture thereof; an anionic surfactant selected from the group consisting of sodium dioctyl sulfosuccinate, sodium dodecylbenzenesulfonate and a mixture thereof; and a fatty acid alkyl ester; said agrochemical and said adjuvant for enhancing the fungicidal efficacy being mixed in a ratio range from 1:0.5 to 1:20 by weight.

20



I

25



II

30

2. The composition of claim 1, wherein the adjuvant for enhancing the fungicidal efficacy is selected from the group consisting of a nonionic surfactant selected from the group consisting of a polyoxyethylene-based nonionic surfactants which has an aliphatic alcohol, a fatty acid or triacyl glyceride as a lipophilic moiety containing at least 8 carbon atoms

and a polyoxyethylene as a hydrophilic moiety having 3 to 25 oxyethylene repeating units; a polyoxyethylene-polyoxypropylene block copolymer containing 2 to 40 oxyethylene and 25 to 45 oxypropylene repeating units and a mixture thereof; an anionic surfactant selected from the group consisting of sodium dioctyl sulfosuccinate, sodium dodecylbenzenesulfonate and a mixture thereof; and fatty acid alkyl esters having at least 14 carbon atoms;

3. The composition of claim 1, which further comprises another agrochemical for preventing or treating plant diseases.

4. An adjuvant composition for enhancing the fungicidal efficacy of an agrochemical of compound of formula (I) or (II) against plant diseases, which comprises;

(a) an effective amount of one or more adjuvants for enhancing the fungicidal efficacy selected from the group consisting of a polyoxyethylene-based nonionic surfactants which has an aliphatic alcohol, a fatty acid or triacyl glyceride as a lipophilic moiety containing at least 8 carbon atoms and a polyoxyethylene as a hydrophilic moiety having 3 to 25 oxyethylene repeating units; a polyoxyethylene-polyoxypropylene block copolymer containing 2 to 40 oxyethylene and 25 to 45 oxypropylene repeating units and a mixture thereof; an anionic surfactant selected from the group consisting of sodium dioctyl sulfosuccinate, sodium dodecylbenzenesulfonate and a mixture thereof; and fatty acid alkyl esters having at least 14 carbon atoms;

- (b) an emulsifier or dispersant; and
- (c) a carrier.

5. A method for preventing or treating a plant disease, which comprises spraying the composition of any one of claims 1 to 5 at a concentration of 4 to 400 mg/l of the agrochemical or 50 to 2,000 mg/l of the adjuvant for enhancing the fungicidal efficacy to said plant.